

REFERENCES

- BLUM, M. S. and J. M. BRAND, (1972) Social insect pheromones: their chemistry and function. Am. Zool. 12: 553.
- KRETSCHMAR, H. C., Z. J. BARNEIS and W. F. ERMAN (1970). The isolation and synthesis of a novel tetracyclic ether from East Indian sandalwood oil. A facile intramolecular Prins reaction. Tet. letters (1), 37.
- MAHESHWARI, M. L., T. C. JAIN, R. B. BATES and S. C. BHATTACHARYYA (1963). Structure and absolute configuration of α -agarofuran, β -agarofuran and dihydroagarofuran. Tetrahedron, 19: 1079.
- MATSUMURA, F., H. C. COPPEL and A. TAI (1968). Isolation and identification of termite trail-following pheromone. Nature, Lond. 219: 963.
- MATSUMURA, F., A. TAI and H. C. COPPEL (1969). Termite trail-following substances, isolation and purification from Reticulitermes virginicus and fungus-infected wood. J. econ. Ent., 62: 599.
- RITTER, F. J. and C. M. A. COENEN-SARABER (1969). Food attractions and a pheromone as trail-following substances for the Saintonge termite. Multiplicity of the trail-following substances in L. trabea-infected wood. Entomologia exp. appl., 12: 611.