

RESEARCH ARTICLE

## Hydrocarbons detection levels in ants

K. Ichinose · A. Lenoir

Received: 4 January 2010 / Revised: 29 April 2010 / Accepted: 3 May 2010  
© International Union for the Study of Social Insects (IUSSI) 2010

**Abstract** In social insects the discrimination process of non-nestmates is generally considered to be based on detection of cuticular hydrocarbons. The required quantity of the stimulus is fundamental but unknown. In laboratory conditions, we demonstrated that *Aphaenogaster senilis* ants are able to discriminate the presence of heterocolonial cuticular hydrocarbons on filter papers at concentrations of  $0.05 \text{ ng/mm}^2$ , the equivalent of  $10^{-4}$  worker. This result does not mean that workers are not able to recognize their own colony odour at lower concentrations, but that the discrimination response appears at very low levels.

**Keywords** Ants · Hydrocarbons detection · *Aphaenogaster*

### Introduction

Social animals discriminate between members of their own colony and conspecific aliens, a process called nestmate recognition. It allows them to maintain colony insularity against competitors and has certainly played an important role in the evolution of eusociality (Wilson and Hölldobler,

2005). Recognition requires the perception of cues emitted by the encountered ant, cues which are compared to the template of the receiver. It is generally thought that cuticular hydrocarbons play an important role in this discrimination (see recent reviews by Hefetz, 2007; d'Ettorre and Lenoir, 2009). A second paradigm involves the role of mandibular gland secretions in nestmate recognition signals (see for example Hernandez et al., 2004), but here we tested the role of non-volatile signals. Nothing is known about the sensibility of the ants to cuticular hydrocarbons. We investigated this question to evaluate the sensibility of workers to a blend of cuticular hydrocarbons.

### Materials and methods

Five colonies of *Aphaenogaster senilis* were collected in the Doñana National Park in Southern Spain. They comprised the queen, numerous brood and 400–800 workers and were maintained in the laboratory at 28°C. They were provided with maggots and a sugared solution for bumblebees three times a week and pieces of orange once a week.

Ten workers were randomly picked up from one nest and killed by freezing. Individual thoraces (plus legs) were dipped in 100 µl pentane for 10 min to extract cuticular hydrocarbons (CHCs). We used only thoraces to remove only cuticular hydrocarbons as the head and gaster have different glands. It was verified that this method removes 90% of the hydrocarbons (A.L., unpublished). The identity of the hydrocarbons was previously determined (Lenoir et al., 2001; Boulay et al., 2007). Different solutions were then prepared from the mother solution (1 ant/10 µl):  $10^{-1}$ ,  $10^{-2}$ ,  $10^{-4}$ ,  $10^{-6}$  and  $10^{-8}$  ant equivalent. To determine the discrimination level of an individual ant towards an

K. Ichinose · A. Lenoir (✉)  
IRBI, UMR CNRS 6035, Institut de Recherche de Biologie de l'Insecte, Faculté des Sciences et Techniques, Université François Rabelais, Parc de Grandmont, 37200 Tours, France  
e-mail: alain.lenoir@univ-tours.fr

K. Ichinose  
e-mail: ichis@affrc.go.jp

Present Address:  
K. Ichinose  
KONARC, Suya 2421, Nishigoshi, Kikuchi, Kumamoto 861-1192, Japan

extract, we used rectangular filter papers ( $5 \times 2$  mm) previously rinsed several times with acetone and pentane to be “odourless.” We verified that these papers contained no hydrocarbons after treatment. The filter paper was impregnated with 10  $\mu\text{l}$  of one of the extracts or with pure pentane. For the test, a worker was introduced in a Petri dish with edges coated with fluon and left for 10 min. The filter paper was deposited and the ant’s behaviour recorded for 2 min. Ants were tested only once with one filter paper. Behaviours were classified according the following categories: antennal contact, opening mandibles, biting and fighting (see Ichinose et al., 2005). Tests were performed with one worker confronted with extracts of the same colony, or to extracts of alien colonies at the different concentrations, or with pure pentane. 1,153 tests were performed.

### Statistics


A principal component analysis was performed on the behavioural data. In a second step, an ANOVA was performed on the first and second components obtained with the PCA analysis using the concentration of extracts and the origin of the extract (pentane, same colony or different colony) as variables.

### Results

The first component of PCA explains 35.78% of the variance and it is correlated with the frequency of aggressions (opening mandibles, biting and fight). An ANOVA indicated that there was a significant difference between homocolonial and heterocolonial tests ( $F_1 = 21.82$ ,  $P = 0.001$ ). The extract concentration did not significantly affect aggression when the worker was confronted with a homocolonial extract ( $F_5 = 0.157$ ,  $P = 0.978$ ) (Fig. 1a). On the contrary, the concentration of the heterocolonial extracts did significantly influence aggression (Fig. 1b;  $F_5 = 3.38$ ,  $P = 0.0048$ ). Aggression was absent or low with pentane. The ants were able to discriminate alien extracts at concentrations as low as  $10^{-4}$  ant but not for  $10^{-6}$  and  $10^{-8}$  ant (post hoc Tukey-HSD test). The second axis of the PCA (21.12% of the variance) was correlated with antennal contacts and the ANOVA confirmed the results of component one (not presented here).

### Discussion

The ant confronted with a worker extract of the same colony did not present the aggressive behaviour which was present when confronted with an alien extract. The detec-



**Fig. 1** First component of the PCA (aggression response)  $\pm$  SE related to the concentration levels of cuticular hydrocarbons (from 1 ant to  $10^{-8}$  ant equivalent). C filter paper impregnated with solvent

tion level for discrimination was very low: at concentrations of  $10^{-4}$  worker equivalent and higher, the workers recognized the alien extract. It has been measured previously that the thorax bears in mean 3,500 ng hydrocarbons (Lenoir et al., 2001), so the worker is sensitive to 0.05 ng/mm<sup>2</sup> on the filter paper, which is very low. The quantity of hydrocarbons on the cuticle related to the surface is not well known and appears to vary considerably with ant species: it is estimated to be from 50 ng/mm<sup>2</sup> in *Cataglyphis iberica* (Dahbi and Lenoir, 1998) to 1,000 ng/mm<sup>2</sup> in *Cataglyphis niger* (Lahav et al., 1998). These strong differences may be due to the calculation methods (see Kroiss et al., 2009), but regardless of the exact value of concentration, our data indicate a discrimination level of  $10^{-4}$  worker equivalent. A similar result has been recently presented on aggressive behaviour of *Camponotus japonicus*: the aggression threshold varying from  $10^{-4}$  to  $10^{-1}$  equivalent of non-nestmate hydrocarbons (Ozaki and Wada-Katsumata, 2010). For other insects only a few data are available: 1,150 ng/mm<sup>2</sup> in the beewolf *Philanthus triangulum* but only 250 ng for the cuckoo wasp parasite *Hedychrum rutilans* which is considered to practice “Chemical insignificance” (Kroiss et al., 2009). For chemical insignificance see Lenoir et al., 1999. The sensitivity of social insects to hydrocarbons is completely unknown. It has been experimentally observed in *Bombus lapidarius* that tricosane (one of the compounds found in the tarsal extracts) at doses  $\geq 10^{-12}$  ng/flower resulted in rejection by foraging workers (Goulson et al., 2000). On the contrary, *Polistes dominulus* react to a foreign founress only when extracts concentrations on a filter paper are higher than two-third wasp equivalent (Cini et al., 2009). The differences are probably due to the experimental designs. One way to estimate the sensitivity of insects to hydrocarbons is to perform electroantennography, but no study has tried to measure it. For example, D’Ettorre et al., 2004 and Ozaki et al., 2005 did not study this question. Batista-Pereira et al., 2004 used a  $10^{-2}$

worker equivalent extract of the termite *Hodotermes tenuis* which released high EAG activity. More experiments are necessary to progress in the comprehension of the detection level of nestmate recognition cues.

**Acknowledgments** We thank Hannah Reynolds (Duke University, NC, USA) who revised the English.

## References

- Batista-Pereira L.G., dos Santos M.G., Correa A.G., Fernandes J.B., Arab A., Costa-Leonardo A.M., Dietrich C., Pereira D.A. and Bueno O.C. 2004. Cuticular hydrocarbons of *Heterotermes tenuis* (Isoptera: Rhinotermitidae): Analyses and electrophysiological studies. *Z. Naturforsch.* **59C**: 135-139
- Boulay R., Hefetz A., Cerdá X., Devers S., Francke W., Twele R. and Lenoir A. 2007. Production of sexuals in a fission-performing ant: dual effects of queen pheromone and colony size. *Behav. Ecol. Sociobiol.* **61**: 1531-1541
- Cini A., Gioli L. and Cervo R. 2009. A quantitative threshold for nestmate recognition in a paper social wasp. *Biol. Lett.* **5**: 459-461
- Dahbi A. and Lenoir A. 1998. Queen and colony odour in the multiple nest ant species, *Cataglyphis iberica* (Formicidae, Formicinae). *Insect. Soc.* **45**: 301-313
- D'Ettorre P., Heinze J., Schulz C., Francke W. and Ayasse M. 2004. Does she smell like a queen? Chemoreception of cuticular hydrocarbon signal in the ant *Pachycondyla inversa*. *J. Exp. Biol.* **207**: 1085-1091
- d'Ettorre P. and Lenoir A. 2009. Nestmate recognition in ants. In: *Ant Ecology* (Lach L., Parr C. and Abbott K., Eds), Oxford, Oxford University Press, pp 190-205
- Goulson D., Stout J.C., Langley J. and Hughes W.O.H. 2000. Identity and function of scent marks deposited by foraging bumblebees. *J. Chem. Ecol.* **26**: 2897-2911
- Hefetz A. 2007. The evolution of hydrocarbon pheromone parsimony in ants (Hymenoptera: Formicidae) - interplay of colony odor uniformity and odor idiosyncrasy. *Myrmecol. News* **10**: 59-68
- Hernandez J.V., Lopez F. and Jaffe K. 2004. Nestmate recognition signals of the leaf-cutting ant *Atta laevigata*. *J. Insect Physiol.* **48**: 287-295
- Ichinose K., Cerdá X., Christides J.-P. and Lenoir A. 2005. Detecting nestmate recognition patterns in the fission-performing ant *Aphaenogaster senilis*: a comparison of different indices. *J. Insect Behav.* **18**: 633-650
- Kroiss J., Schmitt T. and Strohm E. 2009. Low level of cuticular hydrocarbons in a parasitoid of a solitary digger wasp and its potential for concealment. *Entomol. Sci.* **12**: 9-16
- Lahav S., Soroker V., Vander Meer R.K. and Hefetz A. 1998. Nestmate recognition in the ant *Cataglyphis niger*: do queens matter? *Behav. Ecol. Sociobiol.* **43**: 203-212
- Lenoir A., Cuisset D. and Hefetz A. 2001. Effects of social isolation on hydrocarbon pattern and nestmate recognition in the ant *Aphaenogaster senilis* (Hymenoptera: Formicidae). *Insect. Soc.* **48**: 101-109
- Lenoir A., Fresneau D., Errard C. and Hefetz A. 1999. The individuality and the colonial identity in ants: the emergence of the social representation concept. In: *Information Processing in Social Insects* (Detrain C., Deneubourg J.L. and Pasteels J.M., Eds), Basel, Birkhäuser Verlag, pp 219-237
- Ozaki M., Wada-Katsumata A., Fujikawa K., Iwasaki M., Yokohari F., Satoji Y., Nisimura T. and Yamaoka R. 2005. Ant nestmate and non-nestmate discrimination by a chemosensory sensillum. *Science* **309**: 311-314
- Ozaki M. and Wada-Katsumata A. 2010. Perception and olfaction of cuticular compounds. In: *Insect Hydrocarbons. Biology, biochemistry, and chemical ecology* (Blomquist G.J. and Bagnères A.-G., Eds), New York, Cambridge University Press, pp 207-221
- Wilson E.O. and Hölldobler B. 2005. Eusociality: origin and consequences. *Proc. Natl Acad. Sc. USA* **102**: 13367-13371